

Coupling Physical Models to the Epithelial Cell Differentiation

Chris Rycroft¹, Robert Saye¹, Kandice Tanner², Mina Bissel², James Sethian^{1,3}, Daniela Ushizima³,

¹Mathematics Department, University of California, Berkeley, CA;

²Life Sciences Division, Lawrence Berkeley National Lab, Berkeley, CA;

³Computational Research Division, Lawrence Berkeley National Lab, Berkeley, CA.

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

Chris Rycroft¹, Robert Saye¹, Kandice Tanner², Mina Bissel² James
Sethian^{1,3}, Daniela Ushizima³,

¹Mathematics Department, University of California, Berkeley, CA;

²Life Sciences Division, Lawrence Berkeley National Lab, Berkeley, CA;

³Computational Research Division, Lawrence Berkeley National Lab, Berkeley,
CA.

We are making use of techniques from applied mathematics and physics to develop a computational model of a mammalian acinus that is based upon mechanical principles. In our simulation, individual cells are modeled as interacting poroelastic bodies, containing both solid-like and fluid-like phases. We will describe the solid-like phase in detail, where internal stresses can be computed by making use of a reference map, which tracks the amount of deformation of a cell from its initial undeformed state. Each cell has its own deforming boundary; correctly tracking multiple interacting boundaries is a computationally challenging problem, and we will demonstrate some results of recently developed numerical methods that address this. Our model is being compared against two different sets of experiments. First, we are comparing against force measurements of fully grown acini. Second, our results are being compared to three dimensional confocal microscopy images of acini development. As part of this work, we are developing image analysis algorithms to automatically extract features that represent deformations of the cell during the mitotic process. First, we process the hyperstacks using edge-preserving smoothing bilateral filtering, which enhances the contrast between marked structures and background fluorescence. Next, a statistical region merging (SRM) segmentation algorithm is responsible for recovering nuclear mass, for later measurement of volumetric descriptions of the cell. Our framework scales to GB hyperstacks, particularly by splitting the hyperstacks into smaller portions of data, based on the extra-cellular matrix identification, so each subset enclose one acinus. Future work includes identification of consistent segments of similar curvature, aimed at recovering nuclei individually or cell adjacency due to proximity-only or in-mitosis.

This work was partially supported by the Office of Energy Research, U.S. Department of Energy, under Contract Number DE-AC02-05CH11231.